

NELIO®



Nelio® tablets for dogs and cats

Use: For the reduction of proteinuria associated with chronic kidney disease.

Active: Benazepril tailored for veterinary use.

Sizes: Nelio for Dogs 5mg, Nelio for Dogs 20mg, Nelio for Cats 2.5mg, Nelio for Cats 5mg
Packs of 100 tablets.

PRILACTONE®




Prilactone® tablets for dogs

Use: Treats dogs that have congestive heart failure because of faulty heart valves.


Active: Spironolactone

Sizes:

Prilactone Next 10mg (30's)
Prilactone Next 50mg (30's)
Prilactone Next 100mg (24's)




Healthy heart



Valvular heart disease:
The valve is abnormal and loses its tightness. The blood flows back towards the atrium

Dilated cardiomyopathy:
The muscle wall becomes thinner, meaning the heart cannot pump as efficiently. The heart chambers also become larger

Signs of heart failure for dog and cat



interchem at the Heart of Cardiology

Comprehensive Cardiology Product Range



Interchem's Cardiology Academy Series of CPD Cardiology Webinars

For more information contact:
Email: marketing@interchem.ie or ask your local territory manager.

CARDALIS® for the treatment of congestive heart failure caused by chronic degenerative valvular disease in dogs (with diuretic support as appropriate). CARDALIS® contains benazepril and spironolactone. LEGAL CATEGORY: POM IE. AMODIP® for the treatment of systemic hypertension in cats. AMODIP® contains amlodipine. LEGAL CATEGORY: POM IE. ISEMIID® for the treat signs related to congestive heart failure in dogs. ISEMIID® contains torasemide. LEGAL CATEGORY: POM IE. LIBEO® for the treatment of ascites and oedema, particularly associated with cardiac insufficiency in dogs. LIBEO® contains furosemide. LEGAL CATEGORY: POM IE. NELIO® for cats for the reduction of proteinuria associated with chronic kidney disease. NELIO® contains benazepril hydrochloride. LEGAL CATEGORY: POM IE. PRILACTONE® for use in combination with standard therapy for the treatment of congestive heart failure caused by degenerative mitral valve disease in dogs. PRILACTONE® contains spironolactone. LEGAL CATEGORY: POM IE. For information about side effects, precautions, warnings and contraindications always read the product packaging, product leaflet, SPC or datasheet. CARDALIS®, AMODIP®, ISEMIID®, NELIO®, LIBEO® and PRILACTONE® are registered trademarks of Ceva Animal Health, UK. Products distributed in Ireland by Interchem (Ireland) Ltd. Use medicines responsibly.

interchem

Further information is available from:
Interchem (Ireland) Ltd
Unit 29, Cookstown Industrial Estate, Dublin 24
Tel: 01 451 8959 www.interchem.ie

CARDALIS®



Ask your Territory Manager about the Cardalis® App (sleeping respiratory rate)

Cardalis® tablets for dogs

Use: Cardalis is a treatment for heart failure in dogs.


Active: Dual action benazepril and spironolactone.

Cardalis tablets are a combination of Benazepril and Spironolactone which have been compressed in to one handy tablet, helping to manage heart failure which is caused by mitral valve disease in dogs. These tablets should be administered with food, either mixed with a small amount of food offered to the dog just prior to the main meal, or with the meal itself.

Sizes: Small, Medium, Large. Packs of 30's.

Bodyweight (kg) of dog	Small 2.5mg/20mg	Medium 5mg/40mg	Large 10mg/80mg
2.5 - 5	½		
5 - 10	1		
10 - 20		1	
20 - 40			1
40 - 60			1 + ½
60 - 80			2

AMODIP®



Amodip® chewable tablets for cats

Use: Amodip Chewable Tablets are for the treatment of systemic hypertension in cats.

Active: The product contains 1.25 mg amlodipine (as amlodipine besilate) as the active substance. The excipients that are used for the tablet are artificial chicken flavour, malted yeast, microcrystalline cellulose, mannitol, croscarmellose sodium, magnesium stearate and silica colloidal anhydrous. Tablets can be divided into two equal parts to provide flexibility of dosing.

Dosage: Tablets should be administered orally with or without food at a recommended starting dose of 0.125 - 0.25 mg/kg. After 14 days of treatment, the dose may subsequently be doubled or increased up to 0.5 mg /kg once daily if adequate clinical response has not been achieved (e.g. systolic blood pressure remaining over 150 mmHg or a decrease of less than 15 % from the pre-treatment measurement).

Sizes: Pack of 100's.

Bodyweight (kg)	Tablets
2.5-5	½
5.1-10	1
10.1+	2



ISEMID®



Isemid® tablets for dogs

Use: Treats signs related to congestive heart failure in dogs.

Active: Torasemide

Sizes: 1mg, 2mg and 4mg. 9x10's.

LIBEO®



Libeo® chewable tablets for dogs

Use: Treats ascites and oedema, particularly associated with cardiac insufficiency in dogs.

Active: Furosemide

Sizes: 10 mg (120 tablet's), 40mg (120 tablet's).